

# チョコラBBドリンクビットR

Category-3 OTC Drug



## チョコラBB<sup>®</sup>ドリンクビットR

Relief from acne and skin roughness

- Along with vitamins B<sub>2</sub> and B<sub>6</sub>, which are important for skin care, this product contains orotic acid and nicotinamide.
- A pharmaceutical drink that works from inside the body for relief from skin roughness.
- A low-calorie product with 1.2 kcal per bottle.
- It is caffeine-free, so you can take it before bed.

## Indications



Skin roughness



acne or  
pimples



stomatitis

Relief of the following symptoms:

skin roughness, acne/skin eruption, stomatitis, angular stomatitis (swollen or cracked edges of lips), cheilitis (swollen or cracked lips), dermatitis, eczema, rashes, sores, tongue inflammation, hot flushes of the face associated with rosacea, redness of the eye, and itching of the eye. Consult with a physician or pharmacist if no improvement is seen in these symptoms after about 1 month of use.

## Supplementation of vitamin B<sub>2</sub> in the following cases:

When physically fatigued, during pregnancy or lactation, and when physical strength is decreased during and after illness.

## Dosage and Administration

For adults (15 years and over), take 1 bottle (50 mL) once a day.

○ Do not give this drug to children (under 15 years).

## Ingredients

### Ingredients per Bottle (50 mL)

Ingredients	Amount
VitaminB <sub>2</sub> phosphate ester	20 mg
VitaminB <sub>6</sub>	25 mg
VitaminB <sub>1</sub> nitrate	5 mg
Nicotinamide	30 mg
Orotic acid	60 mg

### Additives:

Sodium benzoate, citric acid, sodium citrate, vanillin, paraben, propylene glycol, flavor, acesulfame potassium, erythritol, sucralose, propyl gallate, and DL-malic acid (alcohol 0.1 mL or less).

1. **Ingredients of this product may float or settle out, but this does not affect the efficacy.**
2. **Taking this drug may cause urine to turn yellow. This is due to the absorption of vitamin B2 contained in the drug, with part of it being excreted in the urine. It is not a cause for concern.**



## Precautions

### **⚠ Precautions for Use**

---

#### **Consultation**

1. **If the following symptoms are observed after taking this product, these may be adverse reactions, so immediately discontinue the use of this product and consult a physician, pharmacist, or registered salesperson, being sure to take this product with you.**

Parts Affected	Symptom
Skin	rash
Gastrointestinal system	stomach discomfort

2. **The following symptom may be observed after taking this product. If this symptom persists or worsens, discontinue the use of this product and consult a physician, pharmacist, or registered salesperson, being sure to take this product with you.**

Diarrhea.

3. **When symptoms do not improve even after using the product for about 1 month, discontinue the use of this product and consult a physician, pharmacist, or registered salesperson, being sure to take this product with you.**

#### **Precautions for Storage and Handling**

1. **Store in a cool place away from direct sunlight.**
2. **Keep out of reach of children.**

3. Do not transfer to a different container. (It may lead to misuse or alter the quality of the drug.)

4. Do not use the product after the expiration date.



## Packaging and Price

Packaging	Recommended retail price
50mL	242yen, tax included (220yen, tax excluded)
50mL×3	726yen, tax included (660yen, tax excluded)
50mL×10	2,420yen, tax included (2,200yen, tax excluded)
50mL×50	12,100yen, tax included (11,000yen, tax excluded)
50mL×54	13,068yen, tax included (11,880yen, tax excluded)