

チョコラBBこどもシロップ

Category-3 OTC Drug



チョコラBB[®]こどもシロップ

For decreased physical strength during or after illness, stomatitis, eczema, and tiredness in little children (aged 3 months or over).

Vitamin B₂ is called the “vitamin for fat metabolism” and the “vitamin for skin and mucosa.” Deficiencies of this vitamin can cause symptoms such as fatigue, eczema, dermatitis and stomatitis to occur easily.

Chocora BB Children’s Syrup contains vitamin B₂ as its main ingredient, along with vitamins B₁ and B₆ and biotin, which is known as anti-dermatitis vitamin. It is a peach-flavored syrup for children to swallow easily.

Recommended for improving symptoms such as eczema, acne, dermatitis and stomatitis as well as for supplementation of vitamin B₂ when physical strength is decreased during or after illness in little children aged 3 months or over.



Indications



Skin roughness



acne or
pimples



stomatitis



fatigue and
malaise

Relief of the following symptoms:

Skin roughness, rashes, sores, acne, eczema, dermatitis, stomatitis, angular stomatitis, canker sores, glossitis, rosacea, congestion of the eyes, itchy eyes

Please consult with a physician or pharmacist if no improvement is seen in these symptoms after about 1 month of use.

Supplementation of vitamin B₂ in the following cases:

When physically fatigued or when strength is decreased during and after illness



Dosage and Administration

Measure the following amount with the attached measuring cup and take it twice a day after meals.

Age	Dosage	Number of doses
7 years or over and under 15 years	15 mL	Twice per day
1 year or over and under 7 years	7.5 mL	Twice per day
3 months or over and under 1 year	5mL	Twice per day
Under 3 months	Do not take this medicine	-

Precautions for Dosage and Administration

When allowing children (aged 3 months or over and under 15 years) to take the medicine, a guardian must be present to watch the child and instruct them how to take it.



Ingredients

Ingredients and their actions

One 30 mL bottle contains the following active ingredients.

Ingredients	Amount	Action
Riboflavin sodium phosphate (vitamin B ₂ phosphate ester)	19 mg	<ul style="list-style-type: none">- Supports healthy skin and mucosa.- Aids the metabolism of fats to convert them to energy.
Pyridoxine hydrochloride (vitamin B ₆)	12 mg	<ul style="list-style-type: none">- Supports healthy skin and mucosa.- Aids the metabolism of proteins to convert them to energy.
Thiamine nitrate (vitamin B ₁ nitrate)	6 mg	<ul style="list-style-type: none">- Aids the metabolism of carbohydrates to convert them to energy.
Nicotinamide	20 mg	<ul style="list-style-type: none">- Supports healthy skin and mucosa.
Biotin	250 µg	<ul style="list-style-type: none">- Supports healthy skin and mucosa and suppresses inflammation of the skin.

Additives

Sodium benzoate, ethanol (0.1 mL or less), citric acid, sodium citrate, D-sorbitol, paraben, flavor, hydrogenated maltose starch syrup, glycine, propyl gallate, povidone, DL-malic acid, pH adjuster

Precautions for Ingredients

Taking this medicine may cause urine to turn bright yellow. This is due to the absorption of vitamin B₂ contained in the medicine, with part of it being excreted in the urine. It is not a cause for concern.



Precautions

⚠️ Precautions for Use

Consultation

1. If the following symptoms are observed after taking this drug, these may be adverse reactions. Immediately discontinue the use of this medicine and show this informational leaflet to a physician, pharmacist, or registered salesperson for a consultation.

Parts Affected	Symptom
Skin	Rash
Gastrointestinal system	Stomach discomfort

2. The following symptom may be observed after taking this medicine. If this symptom persists or worsens, discontinue the use of this medicine and show this informational leaflet to a physician, pharmacist, or registered salesperson for a consultation.

Diarrhea

3. If symptoms do not improve even after taking the medicine for about 1 month, stop taking the medicine and consult a physician, pharmacist, or registered salesperson, bringing this informational leaflet with you.

Precautions for storage and handling

1. Store sealed in a cool place away from direct sunlight.
2. Keep out of reach of children.
3. Keep refrigerated after opening the bottle and use it as soon as possible.
4. Do not transfer to a different container. Moreover, do not put other drugs in this container. (It may lead to misuse or alter the quality of the drug.)
5. After taking this drug, rinse the measuring cup and store it securely with the syrup.

6. Do not use the product after the expiration date.



Packaging and Price

Packaging	Recommended retail price
30 mL × 2 (1 box)	770yen, tax included (700yen, tax excluded)
30 mL × 2 × 5 (5 boxes)	3,850yen, tax included (3,500yen, tax excluded)