

本製品は日本国法に基づき製造販売されたものです。
製品および文書は日本国外の法規に準じているわけではありません。
本文書は、日本語の製品情報を翻訳した文書です。使用前に必ずお読みください。
また、必要な時に読めるように大切に保管してください。
This product has been manufactured and sold based on Japanese law.
Neither the product nor this informational leaflet will necessarily conform to the laws of countries outside of Japan.
This leaflet contains a translation of product information from Japanese. You should read this written explanation before using the product.
Store this informational leaflet safely so that it can be read when necessary.



Discovering everything from prevention to cure
Eisai.jp

Category-3 OTC drug

Relief of stiff shoulder, lumbago, and eyestrain

ナボリン®S

Nabolin S

Stiffness of the shoulders and lumbago occur due to excessive loads on the shoulder or low back. If you stay in the same posture or try to maintain a difficult posture, the muscles in your shoulders or low back may harden, damaging the nerves, and causing stiffness or pain.

ナボリン S contains vitamin B₁, which has an effect on muscle fatigue, vitamin E, which improves poor circulation, and activated vitamin B₁₂ (mecobalamin), which repairs peripheral nerve damage. It also contains folic acid, which enhances the action of mecobalamin, which is effective at treating shoulder stiffness and lumbago that feel as if they do not improve when massaged.



Package	Recommended retail price (tax included)
21 tablets	¥1,609
40 tablets	¥2,913
90 tablets	¥4,950
180 tablets	¥9,198

Indications

Relief of the following symptoms:

Muscle and joint pain (shoulder, low back, elbow, and knee pain, stiff shoulder, frozen shoulder, etc.), neuralgia, numbness in the limbs, and eyestrain (chronic eye fatigue and associated blurred vision or pain behind the eye).

Consult with a physician or pharmacist if no improvement is seen in these symptoms after about 1 month of use.

Dosage and Administration

Take the following doses with cold or hot water after a meal.

Age	Dosage	Number of doses
Adult (15 years or older)	1 tablet	3 times a day
Child (Under 15 years)	Do not take this medicine	

Active Ingredients and Action

The daily adult dose of 3 tablets contains the following active ingredients.

Ingredient	Amount	Action
Mecobalamin (activated vitamin B ₁₂)	1500 µg	Promotes the production of components of peripheral nerves (such as proteins and phospholipids) in the body and repairs damaged peripheral nerves.
Folic acid	5 mg	Works with mecobalamin to promote the synthesis of components of peripheral nerves (proteins) in the body and enhances the action of repairing damaged peripheral nerves.
d-α-Tocopherol acetate (natural vitamin E)	100 mg	Maintains the flexibility of red blood cells, which are components of the blood, and blood vessels, improving blood flow.
Fursultiamin hydrochloride (vitamin B ₁ analog)	109.16 mg	Works to change the sugars consumed in food to the energy needed by parts of the body such as muscles, and improves muscle fatigue.
Pyridoxine hydrochloride (vitamin B ₆)	100 mg	Is involved in the metabolism of amino acids and supports the activity of nerves in the body.

Inactive Ingredients

Talc, calcium carbonate, corn starch, lactose, hydroxypropyl cellulose, carnauba wax, calcium silicate, titanium oxide, stearic acid, magnesium stearate, shellac, cellulose, silicon dioxide, white soft sugar, hypromellose, pullulan, povidone, macrogol, dibasic calcium phosphate



Precautions

When not to use the product

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions are more likely to occur.)

This medicine should not be taken in the following persons.

Patients who have had an allergic symptom to this medicine or its ingredients

Consultation

- The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before administration.
Patients undergoing medical treatment from a physician
- If the following symptoms are observed after taking this medicine, these may be adverse reactions, so immediately discontinue the use of this medicine, and show this informational leaflet to a physician, pharmacist, or registered salesperson for a consultation.

Parts affected	Symptom
Skin	Rash, itching
Digestive organs	Nausea and vomiting, stomatitis, loss of appetite, stomach discomfort
Neuropsychiatric system	Dizziness

The following serious symptoms may occur in rare cases. In such cases, immediately seek medical aid:

Name of symptom	Symptom
Shock (anaphylaxis)	Symptoms, such as itching of skin, urticaria, hoarseness, sneezing, itchy throat, breathing difficulties, palpitations, and clouding of consciousness may occur immediately after take.

- The following symptoms may be observed after taking this medicine. If these symptoms persist or worsen, discontinue the use of this medicine, and show this informational leaflet to a physician, pharmacist, or registered salesperson for a consultation.
Constipation, diarrhea, soft stool
- When symptoms do not improve even after taking this medicine for about 1 month, stop taking the medicine and consult a physician, pharmacist or registered salesperson, being sure to take this informational leaflet with you.

Precautions for Storage and Handling

1. Store sealed in the box, in a cool, dry place away from direct sunlight. (This medicine is unstable in light.)
2. Keep out of reach of children.
3. Do not transfer to a different container. Also, do not transfer other medications, etc., into this container. (It may lead to misuse or alter the quality.)
4. A desiccant is included in the container. Do not ingest the desiccant.
5. Do not take tablets that have exceeded their expiration date.
6. From the standpoint of maintaining quality, as a rule of thumb, once the cap has been opened, use the tablets within 6 months, regardless of their expiration date. There is a space marked "Date Opened" on the box's inner flap. Please use that to record the date of opening.
7. When fastening the upper lid of the cap, keep pressing until it clicks.
8. When there are only a few tablets left, it may be difficult to take them out. Remove the cap from the container and take out tablets.

Contact us: For inquiries, please complete the Contact Form. We will endeavor to respond to you as quickly as possible, although there may be some delay depending on the nature of the inquiry and other factors.

Contact Form (English only): https://inquiry.eisai.co.jp/webapp/form/17672_sdab_2/index.do