

サクロンQ

Category-2 OTC Drug



サクロンQ

Saclon Q



Blocks peripheral nerve transmission. Quickly acts on stomach retching, nausea and stomachache.

Saclon Q is a stomach medicine that quickly acts on stomach retching, nausea and stomachache, which are caused by irritation of the stomach mucous membrane by gastric acid, alcohol, or other irritants.

As it is a “quick preparation,” which promptly melts after taking it orally, it is rapidly dissolved in the stomach, so that the ingredient (oxethazaine) directly acts on the stomach mucous membrane to quickly improve stressful symptoms.

You can take it whenever you feel symptoms such as stomachache.

(Do not take this medicine more than 3 times per day. Take the doses at least 4 hours apart.)

Indications



stomachache



hangover

Stomachache, abdominal pain, nausea (retching, stomach retching, retching due to hangovers and overdrinking, sick feeling, and urge to vomit), vomiting, gripping pain (colic, spasms), heartburn, hyperacidity, feeling of discomfort in the stomach, belching (burping)

Dosage and Administration

Take the following dose with cold or hot water.

Age	Dosage	Number of doses
Adults (15 years or over)	2 tablets	Up to 3 times per day
Children (under 15 years)	Do not take this medicine	-

Take the doses at least 4 hours apart.

Precautions for Dosage and Administration

Swallow the tablets quickly without chewing. (These tablets are quickly soluble, so numbness remains in the mouth when the ingredients are spread in the mouth.)

How to take the tablets out

Use your finger and press hard on the raised area of the sheet containing the tablets to break the aluminum foil and remove the tablet. (Accidentally swallowing the tablets while they are still in the sheet can lead to unexpected accidents such as piercing of the esophageal membrane.)

Ingredients

Ingredient and its action

The daily adult dose of 2 tablets contains the following ingredient.

Ingredients	Amount	Action
Oxethazaine	5 mg	Blocks symptoms of irritation, such as stomachache and nausea, by directly acting on the stomach mucous membrane.

Additives

Corn starch, D-mannitol, calcium stearate, cellulose

Precautions

▲ Precautions for Use

When not to use the product

[If you do not follow these instructions, the current symptoms may worsen or adverse reactions are more likely to occur.]

1. This product should not be taken by the following persons:

1. Pregnant women or women who believe they may be pregnant
2. Children aged under 15 years

2. This drug should not be taken together with the following drugs:

Other gastrointestinal drugs (formulations containing scopolia extract), other gastrointestinal pain relief/antispasmodic drugs

3. Do not take this medicine for a long time.

Consultation

1. The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before taking this drug.

1. Patients undergoing medical treatment from a physician
2. The elderly
3. Patients who have experienced allergic symptoms associated with drugs etc.

2. If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug and show this instruction leaflet to your physician, pharmacist, or registered salesperson for a consultation.

Parts Affected	Symptom
Skin	Rash/redness, itching
Mental/nervous system	Headache, drowsiness, dizziness, weakness

3. The following symptom may be observed after taking this drug. If this symptom persists or worsens, discontinue the use of this drug and show this instruction leaflet to your physician, pharmacist, or registered salesperson for a consultation.

Constipation

4. If symptoms do not improve even after taking this medicine 5 to 6 times, stop taking this medicine and consult a physician, pharmacist, or registered salesperson, being sure to take this instruction leaflet with you.

Precautions for storage and handling

1. Store in a cool, low humidity place away from direct sunlight.
2. Keep out of reach of children.
3. Do not transfer to a different container. Moreover, do not put other drugs in this container. (This may lead to misuse or alter the quality of the drug.)
4. Do not use the product after the expiration date.

Packaging and Price

Packaging	Recommended retail price
6 tablets	748yen, tax included (680yen, tax excluded)
12 tablets	1,408yen, tax included (1,280yen, tax excluded)