

saclon q

サクロンQ

Category-2 OTC Drug



<

サクロンブランド情報

製品詳細・添付文書

+

overview

Product Description

サクロンQ



Blocks peripheral nerve transmission. Quickly acts on stomach retching, nausea and stomachache.

Saclon Q is a stomach medicine that quickly acts on stomach retching, nausea and stomachache, which are caused by irritation of the stomach mucous membrane by gastric acid and alcohol, etc. As it is a “quick preparation,” which promptly melts after oral administration, it is rapidly dissolved in the stomach, so that the ingredient (oxethazaine) directly acts on the stomach mucous membrane to quickly improve stressful symptoms.

You can take it whenever you feel symptoms, such as stomachache.

(Do not take this medicine more than 3 times per day, and leave an interval of at least 4 hours between doses.)

Indications

▽ Dosage and Administration

▽ Ingredients

▽ Precautions

▽ Packaging and Price

Indications



Stomachache, abdominal pain, nausea (retching, stomach retching, retching due to hangovers and overdrinking, sick feeling, and feeling of sickness), vomiting, gripping pain (colic, spasms), heartburn, hyperacidity, feeling of discomfort in the stomach, belching (burping)

Dosage and Administration

Take the following dose with cold or hot water.

Age	Dosage	Number of doses
Adults (15 years or over)	2 tablets	Up to 3 times per day
Children (under 15 years)	Do not take this medicine	-

Leave an interval of at least 4 hours between doses.

Precautions for Dosage and Administration

Swallow the tablets quickly without chewing. (These tablets are quickly soluble, so numbness remains in the mouth when the ingredients are spread in the mouth.)

How to Remove the Tablets from the Packaging

Press firmly on a raised part of the sheet containing the tablet to break the aluminum foil on the reverse side, and then take out a tablet and swallow it. (Mistakenly swallowing tablets that are still in their sheet may cause unexpected accidents such as tearing of the esophageal membrane.)

Ingredients

Active Ingredients and Actions

The daily adult dose of 2 tablets contains the following ingredients.

Ingredients	Amount	Action
Oxethazaine	5mg	Blocks symptoms of irritation, such as stomachache, nausea, by directly acting on the stomach mucous membrane.

Additives

Corn starch, D-mannitol, calcium stearate, cellulose

Precautions

⚠ Precautions for Use

When not to use the product

[If you do not follow these instructions, the current symptoms may worsen, or adverse reactions are more likely to occur.]

1. This product should not be taken by the following persons:
 1. Pregnant women or women suspected of being pregnant
 2. Children aged under 15 years
2. This drug should not be taken together with the following drugs: Other gastrointestinal drugs (formulations containing scopolia extract), other pain relief/antispasmodic medicine for the stomach
3. Do not take this medicine for a long time.

Consultation

1. The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before administration.
 1. Patients undergoing medical treatment from a physician
 2. The elderly
 3. Patients who have experienced allergic symptoms associated with drugs, etc.
2. If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this instruction leaflet to your physician, pharmacist, or registered salesperson for a consultation.

Parts Affected	Symptom
Skin	Rash/redness, itching
Neuropsychiatric system	Headache, drowsiness, dizziness, weakness

3. The following symptoms may be observed after taking this drug. If these symptoms persist or worsen, discontinue the use of this drug, and show this instruction leaflet to your physician, pharmacist, or registered salesperson for a consultation.

Constipation

4. When symptoms do not improve even after taking this medicine 5 to 6 times, stop taking this medicine and consult a physician, pharmacist, or registered salesperson, being sure to take this instruction leaflet with you.

Precautions for storage and handling

1. Store in a cool, low humidity place away from direct sunlight.
2. Keep out of reach of children.
3. Do not transfer to a different container. Moreover, do not put other drugs in this container. (This may lead to misuse or alter the quality of the drug.)
4. Do not use the product after the expiration date.

Packaging and Price

Packaging	Recommended retail price
6 tablets	748yen, tax included (680yen, tax excluded)
12 tablets	1,408yen, tax included (1,280yen, tax excluded)