

トラベルミン チュロップ レモン味 (Brand name : トラベルミン チュロップ レモン味)

Category-2 OTC Drug



トラベルミン[®] チュロップ レモン味

For prevention and relief of motion sickness

Brandname : トラベルミン チュロップレモン味

Travelmin Churopp Lemon Flavor is a medicine that prevents and relieves symptoms associated with motion sickness, such as dizziness, nausea, and headache, to help you to comfortably enjoy your trip and going out.

It is a lozenge-type medicine which is easy to take for children. You can easily take it on the spot when you're in a hurry to go out or when you're feeling sick.



Indications



Motion
sickness

Prevention and relief of dizziness, nausea, and headache associated with motion sickness



Dosage and Administration

To prevent motion sickness, take the following dose 30 minutes before getting in a vehicle, either by chewing the medicine or dissolving it in your mouth.

Age	Dosage	Number of doses
15 years or over	2 tablets	Up to 2 times in intervals of at least 4 hours
11 to 14 years	2 tablets	Up to 2 times in intervals of at least 4 hours
5才~10才	1 tablets	Up to 2 times in intervals of at least 4 hours
Less than 5 years	Do not take this medicine	-

When taking additional doses, take the doses at least 4 hours apart. Do not take this medicine more than twice per day.

Precautions for Dosage and Administration

When allowing children (5 to 14 years) to take the medicine, a guardian must be present to watch the child and instruct them how to take it. When giving the child the medicine to be taken while out of the house, such as while on a school trip, give them instructions about the dosage, how to take the medicine, etc.

Note

- This medicine is taken either by chewing it or by dissolving it in your mouth. If it is difficult to chew, dissolve it before taking it so that you do not damage your teeth. Take care that it does not accidentally become stuck in your throat.
- Take 1 tablet at a time. Do not put 2 tablets in your mouth at once.



Ingredients

Ingredients and their actions

The dose of 2 tablets (11 years or over) contains the following ingredients.

Ingredients	Amount	Action
d-Chlorpheniramine maleate	1.33 mg	Acts on the autonomic nerves, preventing and relieving the symptoms of motion sickness.
Scopolamine hydrobromide hydrate	0.166 mg	Reduces sensory confusion while riding in a vehicle.

Additives

Citric acid, l-menthol, flavor, acesulfame potassium, isomalt, glycerol esters of fatty acids, vitamin B₂



Precautions

⚠️ Precautions for Use

When not to use the product

If you do not follow these instructions, the current symptoms may worsen or adverse reactions/incidents are more likely to occur.

1. This drug should not be taken together with the following drugs:

Other medicines for motion sickness, cold medicines, antipyretics/analgesics, sedatives, expectorants, medicines for indigestion and heartburn relief, medicines taken orally that contain antihistamines, etc.

(such as medicines taken orally for rhinitis or medicines for allergies)

2. Do not operate vehicles or machinery after taking this drug. (Symptoms such as sleepiness, blurred vision, and increased sensitivity to light may occur.)

Consultation

1. The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before taking this medicine.
 1. Patients undergoing medical treatment from a physician
 2. Pregnant women or women who believe they may be pregnant
 3. The elderly
 4. Patients who have experienced allergic symptoms associated with drugs etc.
 5. Persons with difficulty urinating
 6. Persons diagnosed with glaucoma or heart disease
2. If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug and show this box to your physician, pharmacist, or registered salesperson for a consultation.

Parts Affected	Symptom
Skin	Rash/redness, itching
Mental/nervous system	Headache
Urinary system	Difficulty urinating
Other	Facial hot flashes, increased sensitivity to light

On rare occasions, the following serious symptoms may occur. In this case, visit a physician immediately.

Name of symptom	Symptom
Aplastic anemia	Symptoms such as bruises, nose bleeds, bleeding gums, fever, pale appearance of skin and mucosa, fatigue, palpitations, shortness of breath, becoming dizzy from feeling sick and blood in the urine may occur.

Agranulocytosis	Symptoms such as acute high fever, chills and sore throat may occur.
-----------------	--

3. The following symptoms may be observed after taking this medicine. If these symptoms persist or worsen, discontinue the use of this drug and show this box to your physician, pharmacist, or registered salesperson for a consultation.

Dry mouth, constipation, drowsiness, blurred vision

Precautions for storage and handling

1. Store in a cool, low humidity place away from direct sunlight.
2. Keep out of reach of children.
3. Do not transfer to a different container. (This may lead to misuse or alter the quality of the drug.)
4. Do not use the product after the expiration date.



Packaging and Price

Packaging	Recommended retail price
6 tablets	521 yen, tax included (474 yen, tax excluded)