

travelmin

# トラベルミン（大人用）

Category-2 OTC Drug



< [トラベルミンブランド情報](#)

[製品詳細・添付文書](#) +

overview

## Product Description

### トラベルミン®

#### For prevention and relief of motion sickness

Travelmin is a motion sickness medicine for adults that effectively prevents and relieves the symptoms of motion sickness.

If you are worried about feeling sick, you can prevent motion sickness symptoms by taking it 30 minutes before you board the vehicle.

Even after you feel sick, you can take it to improve the symptoms of motion sickness (dizziness, nausea, and headaches), helping you to enjoy your trip.

Indications

▽ Dosage and Administration

▽ Ingredients

▽ Precautions

▽ Packaging and Price

### Indications



Prevention and relief of dizziness, nausea, and headache associated with motion sickness

## Dosage and Administration

To prevent motion sickness, take the following dose 30 minutes before boarding, either by chewing the medicine or dissolving it in your mouth.

Age	Dosage	Number of doses
15 years or over	1tablet	Up to 3 times in intervals of at least 4 hours
Less than 15 years	Do not take this medicine	-

When taking additional doses, leave an interval of at least 4 hours between doses. Do not take this medicine more than 3 times per day.

### How to take the tablets out

As shown in the figure below, press hard on the raised area of the sheet containing the tablets with a finger to break the aluminum foil and then remove and take the tablet. (Accidentally swallowing the tablets while they are still in the sheet can lead to unexpected accidents such as piercing of the esophageal membrane.)



## Ingredients

### Ingredients and their actions

One tablet contains the following ingredients.

Ingredients	Amount	Action
Diphenhydramine salicylate	40mg	Acts on the autonomic nerves, preventing and relieving the symptoms of motion sickness.
Diprophylline	26mg	Suppresses sensory confusion that occurs due to swaying motions, preventing motion sickness.

### Additives

Talc, corn starch, lactose, carmellose calcium, hydrogenated oil, titanium oxide, calcium stearate, cellulose, hypromellose, macrogol, silicon dioxide

## Precautions

## ▲ Precautions for Use

### When not to use the product

If you do not follow these instructions, the current symptoms may worsen or adverse reactions/incidents are more likely to occur.

1. This product should not be taken by the following persons:

Persons diagnosed as having the following  
Glaucoma, prostatic hypertrophy

2. This drug should not be taken together with the following drugs:

Other medicine for motion sickness, cold medicine, antipyretics/analgesics, sedatives, expectorant, internal medicine containing antihistamines, etc. (such as internal medicine for rhinitis and medicine for allergies)

3. Do not operate vehicles or machinery after taking this drug.  
(Sleepiness may occur.)
4. Nursing women should either not take this medicine or stop nursing while taking this medicine.

### Consultation

1. The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before administration.
  1. Patients undergoing medical treatment from a physician
  2. Pregnant women or women suspected of being pregnant
  3. Patients who have experienced allergic symptoms associated with drugs, etc.
  4. Persons with the following symptoms:  
Urination difficulty
  5. Persons diagnosed as having the following:  
Epilepsy, thyroid disease
2. If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this instruction leaflet to your physician, pharmacist, or registered salesperson for a consultation.

Parts Affected	Symptom
Skin	Rash/redness, itching
Cardiovascular system	Palpitations
Urinary system	Urination difficulty

3. The following symptoms may be observed after taking this drug. If these symptoms persist or worsen, discontinue the use of this drug, and show this instruction leaflet to your physician, pharmacist, or registered salesperson for a consultation.

Dry mouth, drowsiness

### Other precautions

Note that the action of this drug may become stronger if you drink alcohol while taking it.

### Precautions for storage and handling

1. Store in a cool, low humidity place away from direct sunlight.
2. Keep out of reach of children.
3. Do not transfer to a different container. Moreover, do not put other drugs in this container.  
(This may lead to misuse or alter the quality of the drug.)
4. Do not use the product after the expiration date.

## Packaging and Price

Packaging	Recommended retail price
6 tablets	660yen, tax included (600yen, tax excluded)